

Some New Generalized Trigonometric, Hyperbolic and Exponential Fuzzy Entropy and Fuzzy Directed Divergence Measures

Maheshwari^{1,*}, Sapna Thakur² and M.A. Siddiqui³

^{1,3}*Department of Mathematics, Govt. V. Y. T. P. G. Autonomous College, Durg, Chhattisgarh*

²*Department of Mathematics, Bhilai Mahila Mahavidyalaya, Bhilai, Chhattisgarh*

Abstract

In the existing literature of information theory, there is a huge availability of measures of fuzzy entropies each with its own merits and limitations. Keeping in view the flexibility in the system and application areas and by using the concept of fuzzy set we have introduced some new generalized trigonometric, hyperbolic and exponential measures of fuzzy entropy and fuzzy directed divergence.

Keywords: Fuzzy Entropy, Fuzzy Directed Divergence, Measures of Fuzzy Information.

1. Introduction

Uncertainty and fuzziness are the basic nature of human thinking and of many real world objectives. Fuzziness is found in our decision, in our language and in the way we process information. The main use of information is to remove uncertainty and fuzziness. In fact, we measure information supplied by the amount of probabilistic uncertainty removed in an experiment and the measure of uncertainty removed is also called as a measure of information while measure of fuzziness is the measure of vagueness and ambiguity of uncertainties. Shannon [2] used “entropy” to measure uncertain degree of the randomness in a probability distribution. Let X is a discrete random variable with probability distribution $P = (p_1, p_2, \dots, p_n)$ in an experiment. The information contained in this experiment is given by

$$H(P) = - \sum_{i=1}^n p_i \log p_i \quad (1)$$

Which is well known Shannon entropy.

The concept of entropy has been widely used in different areas, e.g. communication theory, statistical mechanics, finance, pattern recognition, and neural network etc. Fuzzy set theory developed by Lofti A. Zadeh [8] has found wide applications in many areas of science and technology, e.g. clustering, image processing, decision making etc. because of its capability to model non-statistical imprecision or vague concepts.

It may be recalled that a fuzzy subset A in U (universe of discourse) is characterized by a *membership function* $\mu_A: U \rightarrow [0,1]$ which represents the *grade of membership* of $x \in U$ in A as follows

$$\begin{aligned}\mu_A(x) &= 0 \text{ if } x \text{ does not belongs to } A, \\ &\quad \text{and there is no uncertainty} \\ &= 1 \text{ if } x \text{ belongs to } A \text{ and there is no uncertainty} \\ &= 0.5 \text{ if maximum uncertainty}\end{aligned}$$

In fact $\mu_A(x)$ associates with each $x \in U$ a grade of membership in the set A . When $\mu_A(x)$ is valued in $\{0,1\}$ it is the characteristic function of a crisp (i.e. nonfuzzy) set. Since $\mu_A(x)$ and $1 - \mu_A(x)$ gives the same degree of fuzziness, therefore, corresponding to the entropy due to Shannon [2], De Luca and Termini [1] suggested the following measure of fuzzy entropy:

$$H(A) = -[\sum_{i=1}^n \mu_A(x_i) \log \mu_A(x_i) + \sum_{i=1}^n (1 - \mu_A(x_i)) \log (1 - \mu_A(x_i))] \quad (2)$$

De Luca and Termini introduced a set of properties and these properties are widely accepted as a criterion for defining any new fuzzy entropy. In fuzzy set theory, the entropy is a measure of fuzziness which expresses the amount of average ambiguity/difficulty in making a decision whether an element

belongs to a set or not. So, a measure of average fuzziness in a fuzzy set should have at least the following properties to be valid fuzzy entropy:

- i) $H(A) = 0$ when $\mu_A(x_i) = 0$ or 1 .
- ii) $H(A)$ increases as $\mu_A(x_i)$ increases from 0 to 0.5 .
- iii) $H(A)$ decreases as $\mu_A(x_i)$ increases from 0.5 to 1 .
- iv) $H(A) = H(\bar{A})$, i.e. $\mu_A(x_i) = 1 - \mu_A(x_i)$
- v) $H(A)$ is a concave function of $\mu_A(x_i)$.

Kullback and Leibler [7] obtained the measure of directed divergence of probability distribution

$P = (p_1, p_2, \dots, p_n)$ from the probability distribution $Q = (q_1, q_2, \dots, q_n)$ as

$$D(P:Q) = \sum_{i=1}^n p_i \log \frac{p_i}{q_i} \quad (3)$$

Let A and B be two standard fuzzy sets with same supporting points x_1, x_2, \dots, x_n and with fuzzy vectors $\mu_A(x_1), \mu_A(x_2), \dots, \mu_A(x_n)$ and $\mu_B(x_1), \mu_B(x_2), \dots, \mu_B(x_n)$. The simplest measure of fuzzy directed divergence as suggested by Bhandari and Pal (1993), is

$$D(A:B) = \sum_{i=1}^n \mu_A(x_i) \log \frac{\mu_A(x_i)}{\mu_B(x_i)} + \sum_{i=1}^n (1 - \mu_A(x_i)) \log \frac{(1 - \mu_A(x_i))}{(1 - \mu_B(x_i))} \quad (4)$$

satisfying the conditions:

- i) $D(A:B) \geq 0$
- ii) $D(A:B) = 0$ iff $A = B$
- iii) $D(A:B) = D(B:A)$
- iv) $D(A:B)$ is a convex function of $\mu_A(x_i)$

later kapur [5],[6] introduced a number of trigonometric hyperbolic and exponential measures of fuzzy entropy and fuzzy directed divergence. In section 2 and 3 we introduce some new trigonometric, hyperbolic and exponential measures of fuzzy entropy and measures of fuzzy directed divergence.

2. New Measures of Fuzzy Entropy

2.1 Trigonometric Measure of Fuzzy Entropy

Consider the function $\sin \pi x$ where $0 \leq x \leq 1$, is a convex function which gives us

$$H_1(A) = \sum_{i=1}^n \sin \left(\frac{\pi}{\mu_A(x_i)} \right) + \sum_{i=1}^n \sin \left(\frac{\pi}{(1 - \mu_A(x_i))} \right) \quad (5)$$

is a new measure of fuzzy entropy.

in particular for $\beta \leq \pi$

$$H_2(A) = \sum_{i=1}^n \sin \left(\frac{\beta}{\mu_A(x_i)} \right) + \sum_{i=1}^n \sin \left(\frac{\beta}{(1 - \mu_A(x_i))} \right) - \sin \beta \quad (6)$$

is also a new measure of fuzzy entropy.

(5) is a special case of (6) when $\beta = \pi$.

Another special case of (6) arises when $\beta = \frac{\pi}{2}$ we get

$$H_3(A) = H_2(A) = \sum_{i=1}^n \sin \left(\frac{\pi}{2\mu_A(x_i)} \right) + \sum_{i=1}^n \sin \left(\frac{\pi}{2(1 - \mu_A(x_i))} \right) - 1 \quad (7)$$

Another trigonometric measure of fuzzy entropy is

$$H_4(A) = \sum_{i=1}^n \sin\left(\frac{\beta}{\mu_A(x_i)} + \alpha\right) + \sum_{i=1}^n \sin\left(\frac{\beta}{(1-\mu_A(x_i))} + \alpha\right) - \sin(\beta + \alpha) \quad (8)$$

(8) reduces to (6) when $\alpha = 0$.

(8) reduces to (7) when $\alpha = 0, \beta = \frac{\pi}{2}$.

(8) reduces to (5) when $\alpha = 0, \beta = \pi$.

(8) is a 2-parameter measure of fuzzy entropy.

If we put $\alpha = \frac{\pi}{2}$ we get

$$H_5(A) = \sum_{i=1}^n \cos\left(\frac{\beta}{\mu_A(x_i)}\right) + \sum_{i=1}^n \cos\left(\frac{\beta}{(1-\mu_A(x_i))}\right) - \cos\beta \quad (9)$$

is a new measure of fuzzy entropy. Clearly above given measures of fuzzy entropy are satisfying all the properties which are given in section 1. So these are valid measures of fuzzy entropy.

2.2 Hyperbolic Measure of Fuzzy Entropy

\sinhx, \coshx, \tanhx where $0 \leq x \leq 1$ are all convex functions and gives us following valid measures of fuzzy entropy

$$H_6(A) = \sinh\beta - \sum_{i=1}^n \sinh\left(\frac{\beta}{\mu_A(x_i)}\right) - \sum_{i=1}^n \sinh\left(\frac{\beta}{(1-\mu_A(x_i))}\right) \quad (10)$$

$$H_7(A) = \cosh\beta - \sum_{i=1}^n \cosh\left(\frac{\beta}{\mu_A(x_i)}\right) - \sum_{i=1}^n \cosh\left(\frac{\beta}{(1-\mu_A(x_i))}\right) \quad (11)$$

$$H_8(A) = \tanh\beta - \sum_{i=1}^n \tanh\left(\frac{\beta}{\mu_A(x_i)}\right) - \sum_{i=1}^n \tanh\left(\frac{\beta}{(1-\mu_A(x_i))}\right) \quad (12)$$

Since $x^m \sinhx, x^m \coshx, x^m \tanhx$ are also convex functions for $m \geq 1$, we get the following additional measures of fuzzy entropy.

$$H_9(A) = \sinh\beta - \sum_{i=1}^n \frac{1}{\mu_A^m(x_i)} \sinh\left(\frac{\beta}{\mu_A(x_i)}\right) - \sum_{i=1}^n \frac{1}{(1-\mu_A(x_i))^m} \sinh\left(\frac{\beta}{(1-\mu_A(x_i))}\right) \quad (13)$$

$$H_{10}(A) = \cosh\beta - \sum_{i=1}^n \frac{1}{\mu_A^m(x_i)} \cosh\left(\frac{\beta}{\mu_A(x_i)}\right) - \sum_{i=1}^n \frac{1}{(1-\mu_A(x_i))^m} \cosh\left(\frac{\beta}{(1-\mu_A(x_i))}\right) \quad (14)$$

$$H_{11}(A) = \tanh\beta - \sum_{i=1}^n \frac{1}{\mu_A^m(x_i)} \tanh\left(\frac{\beta}{\mu_A(x_i)}\right) - \sum_{i=1}^n \frac{1}{(1-\mu_A(x_i))^m} \tanh\left(\frac{\beta}{(1-\mu_A(x_i))}\right) \quad (15)$$

2.3 Exponential Measures of Fuzzy Entropy

Since $x^m e^{ax}$ is a convex function when $m \geq 1, x > 0$ we get the measure of fuzzy entropy

$$H_{12}(A) = e^a - \sum_{i=1}^n \frac{1}{\mu_A^m(x_i)} e^{\frac{a}{\mu_A(x_i)}} - \sum_{i=1}^n \frac{1}{(1-\mu_A(x_i))^m} e^{\frac{a}{(1-\mu_A(x_i))}} \quad (16)$$

3. New Measures of Fuzzy Directed Divergence

3.1 New Hyperbolic Measures of Fuzzy Directed Divergence

Using the convexity of $\sinh x, \cosh x, \tanh x$ we get the following measures of hyperbolic fuzzy directed divergence.

$$D_1(A: B) = \sum_{i=1}^n \frac{1}{\mu_B(x_i)} \sinh \left(\beta \frac{\mu_B(x_i)}{\mu_A(x_i)} \right) + \sum_{i=1}^n \left(\frac{1}{1 - \mu_B(x_i)} \right) \sinh \left(\beta \frac{(1 - \mu_B(x_i))}{(1 - \mu_A(x_i))} \right) - \frac{1}{\mu_B(x_i)} \sinh \beta - \left(\frac{1}{1 - \mu_B(x_i)} \right) \sinh \beta \quad (17)$$

$$D_2(A: B) = \sum_{i=1}^n \frac{1}{\mu_B(x_i)} \cosh \left(\beta \frac{\mu_B(x_i)}{\mu_A(x_i)} \right) + \sum_{i=1}^n \left(\frac{1}{1 - \mu_B(x_i)} \right) \cosh \left(\beta \frac{(1 - \mu_B(x_i))}{(1 - \mu_A(x_i))} \right) - \frac{1}{\mu_B(x_i)} \cosh \beta - \left(\frac{1}{1 - \mu_B(x_i)} \right) \cosh \beta \quad (18)$$

$$D_3(A: B) = \sum_{i=1}^n \frac{1}{\mu_B(x_i)} \tanh \left(\beta \frac{\mu_B(x_i)}{\mu_A(x_i)} \right) + \sum_{i=1}^n \left(\frac{1}{1 - \mu_B(x_i)} \right) \tanh \left(\beta \frac{(1 - \mu_B(x_i))}{(1 - \mu_A(x_i))} \right) - \frac{1}{\mu_B(x_i)} \tanh \beta - \left(\frac{1}{1 - \mu_B(x_i)} \right) \tanh \beta \quad (19)$$

Again since $x^m \sinh x, x^m \cosh x, x^m \tanh x$ are also convex functions for $m \geq 1$, we get the following more general hyperbolic measures of fuzzy directed divergence.

$$D_4(A: B) = \sum_{i=1}^n \frac{1}{\mu_A^m(x_i) \mu_B^{1-m}(x_i)} \sinh \left(\beta \frac{\mu_B(x_i)}{\mu_A(x_i)} \right) + \sum_{i=1}^n \frac{1}{(1 - \mu_A(x_i))^m (1 - \mu_B(x_i))^{1-m}} \sinh \left(\beta \frac{(1 - \mu_B(x_i))}{(1 - \mu_A(x_i))} \right) - \frac{1}{\mu_B(x_i)} \sinh \beta - \frac{1}{(1 - \mu_B(x_i))} \sinh \beta \quad (20)$$

$$D_5(A: B) = \sum_{i=1}^n \frac{1}{\mu_A^m(x_i) \mu_B^{1-m}(x_i)} \cosh \left(\beta \frac{\mu_B(x_i)}{\mu_A(x_i)} \right) + \sum_{i=1}^n \frac{1}{(1 - \mu_A(x_i))^m (1 - \mu_B(x_i))^{1-m}} \cosh \left(\beta \frac{(1 - \mu_B(x_i))}{(1 - \mu_A(x_i))} \right) - \frac{1}{\mu_B(x_i)} \cosh \beta - \frac{1}{(1 - \mu_B(x_i))} \cosh \beta \quad (21)$$

$$\begin{aligned}
D_6(A:B) &= \sum_{i=1}^n \frac{1}{\mu_A^m(x_i) \mu_B^{1-m}(x_i)} \tanh \left(\beta \frac{\mu_B(x_i)}{\mu_A(x_i)} \right) \\
&+ \sum_{i=1}^n \frac{1}{(1-\mu_A(x_i))^m (1-\mu_B(x_i))^{1-m}} \\
&\tanh \left(\beta \frac{(1-\mu_B(x_i))}{(1-\mu_A(x_i))} \right) - \frac{1}{\mu_B(x_i)} \tanh \beta - \frac{1}{(1-\mu_B(x_i))} \tanh \beta
\end{aligned} \tag{22}$$

3.2 New Exponential Measures of Fuzzy Directed Divergence

Since $x^m e^{ax}$ is a convex function when $m \geq 1, x > 0$ we get the following measures of fuzzy directed divergence

$$\begin{aligned}
D_7(A:B) &= \sum_{i=1}^n \frac{1}{\mu_A^m(x_i) \mu_B^{1-m}(x_i)} e^{a \left(\frac{\mu_B(x_i)}{\mu_A(x_i)} \right)} \\
&+ \sum_{i=1}^n \frac{1}{(1-\mu_A(x_i))^m (1-\mu_B(x_i))^{1-m}} e^{a \left(\frac{(1-\mu_B(x_i))}{(1-\mu_A(x_i))} \right)} - e^a
\end{aligned} \tag{23}$$

Special case for $m=0$ and $m=1$ are

$$D_8(A:B) = \sum_{i=1}^n \frac{1}{\mu_B(x_i)} e^{a \left(\frac{\mu_B(x_i)}{\mu_A(x_i)} \right)} + \sum_{i=1}^n \frac{1}{(1-\mu_B(x_i))} e^{a \left(\frac{(1-\mu_B(x_i))}{(1-\mu_A(x_i))} \right)} - e^a \tag{24}$$

$$D_9(A:B) = \sum_{i=1}^n \frac{1}{\mu_A(x_i)} e^{a \left(\frac{\mu_B(x_i)}{\mu_A(x_i)} \right)} + \sum_{i=1}^n \frac{1}{(1-\mu_A(x_i))} e^{a \left(\frac{(1-\mu_A(x_i))}{(1-\mu_B(x_i))} \right)} - e^a \tag{25}$$

4. Conclusion

In section 2 and 3 by using the convexity of some trigonometric, hyperbolic and exponential function and satisfying the conditions of fuzzy entropy and fuzzy directed divergence we get some new trigonometric, hyperbolic and exponential measures of fuzzy entropy and fuzzy directed divergence.

5. References

- [1] A. De Luca and S. Termini, *A Definition of a Non-probabilistic Entropy in the Setting of fuzzy sets theory*, Information and Control, 20, 301-312, 1972.
- [2] C.E.Shannon (1948). "A Mathematical Theory of Communication". Bell. System Tech. Journal Vol. 27, pp. 379-423,623-659.
- [3] D. Bhandari and N. R. Pal, *Some new information measures for fuzzy sets*, Information Science, 67, 204 - 228, 1993.

- [4] F.M.Reza (1948&1949). "An introduction to information theory". Mc. Graw-Hill, New-York.
- [5] J.N.Kapur. Some New Measures of Directed Divergence. Willey Eastern Limited.
- [6] J.N.Kapur. Trigonometrical Hyperbolic and Exponential Measure of Fuzzy Entropy and Fuzzy Directed Divergence. MSTS
- [7] S. Kullback and R.A. Leibler, *On Information and Sufficiency*, Annals of Mathematical Statistics, 22,79 - 86, 1951.
- [8] L. A. Zadeh, *Fuzzy Sets*, Information and Control, 8, 338 - 353, 1965.