

































- status", *New Phytol.*, 159, pp. 213–228. doi:10.1046/j.1469-8137.2003.00790.x.
- [39] Manetas, Y., Petropoulou, Y., Psaras, G. K., and Drinia, A., 2003, "Exposed red (anthocyanic) leaves of *Quercus coccifera* display shade characteristics", *Funct. Plant Biol.*, 30, pp. 265–270. doi:10.1071/FP02226.
- [40] Essaghi, S., Hachmi, M., Yessef, M., and Dehhaoui, M., 2016, "Leaf shrinkage : a predictive indicator of the potential variation of the surface area - to - volume ratio according to the leaf moisture content", *Springerplus*, 5(1). doi:10.1186/s40064-016-2900-3.
- [41] Hulshof, C. M., and Swenson, N. G., 2010, "Variation in leaf functional trait values within and across individuals and species: An example from a Costa Rican dry forest", *Funct. Ecol.*, 24, pp. 217–223. doi:10.1111/j.1365-2435.2009.01614.x.
- [42] Barazani, O., Dudai, N., and Golan-Goldhirsh, A., 2003, "Comparison of mediterranean *Pistacia lentiscus* genotypes by random amplified polymorphic DNA, chemical, and morphological analyses", *J. Chem. Ecol.*, 29, pp. 1939–1952. doi:10.1023/A:1024862614345.
- [43] Werner, C., Correia, O., and Beyschlag, W., 1999, "Two different strategies of Mediterranean macchia plants to avoid photoinhibitory damage by excessive radiation levels during summer drought", *Acta Oecologica*, 20, pp. 15–23. doi:10.1016/S1146-609X(99)80011-3.
- [44] Castro, F. X., Tudela, A., and Sebastià, M. T., 2003, "Modeling moisture content in shrubs to predict fire risk in Catalonia (Spain)", *Agric. For. Meteorol.*, 116, pp. 49–59. doi:10.1016/S0168-1923(02)00248-4.
- [45] Chaves, J., Maroco, M. M., and Pereira, J. P., 2003, "Understanding plant responses to drought — from genes to whole plant", *Func Plant Biol Review* :, *Funct. Plant Biol.*, 30, pp. 239–264. doi:10.1071/FP02076.
- [46] Aranda, I., Ramírez-Valiente, J. A., and Rodríguez-Calcerrada, J., 2014, "Características funcionales que influyen en la respuesta a la sequía de las especies del género *Quercus*: variación inter- e intra-específica", *Ecosistemas*, 23, pp. 27–36.
- [47] Sánchez-Gómez, D., Valladares, F., and Zavala, M. A., 2006, "Performance of seedlings of Mediterranean woody species under experimental gradients of irradiance and water availability: Trade-offs and evidence for niche differentiation", *New Phytol.*, 170, pp. 795–806. doi:10.1111/j.1469-8137.2006.01711.x.
- [48] Gratani, L., Ghia, E., 2002, "Adaptive strategy at the leaf level of *Arbutus unedo* L. to cope with Mediterranean climate", *Flora - Morphol. Distrib. Funct. Ecol. Plants*, 197, pp. 275–284. doi:10.1078/0367-2530-00041.