# **Bounds of the Mertens Functions**

## Darrell Cox<sup>a</sup>, Sourangshu Ghosh<sup>b</sup>, Eldar Sultanow<sup>c</sup>

<sup>a</sup>Grayson County College, Denison, TX 75020, USA

<sup>b</sup>Department of Civil Engineering , Indian Institute of Technology Kharagpur, West Bengal, India

<sup>c</sup>Potsdam University, 14482 Potsdam, Germany

### **Abstract**

In this paper we derive new properties of Mertens function and discuss about a likely upper bound of the absolute value of the Mertens function  $\sqrt{\log(x!)} > |M(x)|$  when x > 1. Using this likely bound we show that we have a sufficient condition to prove the Riemann Hypothesis.

#### 1. INTRODUCTION

We define the Mobius Function  $\mu(k)$ . Depending on the factorization of n into prime factors the function can take various values in  $\{-1, 0, 1\}$ 

- $\mu(n) = 1$  if n has even number of prime factors and it is also square-free(divisible by no perfect square other than 1)
- $\mu(n) = -1$  if n has odd number of prime factors and it is also square-free
- $\mu(n) = 0$  if n is divisible by a perfect square.

Mertens function is defined as  $M(n) = \sum_{k=1}^{n} \mu(k)$  where  $\mu(k)$  is the Mobius function. It can be restated as the difference in the number of square-free integers up to x that have even number of prime factors and the number of square-free integers up to x that have odd number of prime factors. The Mertens function rather grows very slowly since the Mobius function takes only the value  $0, \pm 1$  in both the positive and

negative directions and keeps oscillating in a chaotic manner. Mertens after verifying all the numbers up to 10,000 conjectured that the absolute value of M(x) is always bounded by  $\sqrt{x}$ . This conjecture was later disproved by Odlyzko and te Riele<sup>1</sup>. This conjecture is replaced by a weaker one by Stieltjes<sup>2</sup> who conjectured that  $M(x) = O(x^{\frac{1}{2}})$ . Littlewood<sup>3</sup> proved that the Riemann hypothesis is equivalent to the statement that for every  $\epsilon > 0$  the function  $M(x)x^{-\frac{1}{2}-\epsilon}$  approaches zero as  $x \to \infty$ . This proves that the Riemann Hypothesis is equivalent to conjecture that  $M(x) = O(x^{\frac{1}{2}+\epsilon})$  which gives a rather very strong upper bound to the growth of M(x). Although there exists no analytic formula, Titchmarsh<sup>4</sup> showed that if the Riemann Hypothesis is true and if there exist no multiple non-trivial Riemann zeta function zeros, then there must exist a sequence  $T_k$  which satisfies  $k \le T_k \le k+1$  such that the following result holds:

$$M_0(x) = \lim_{k \to \infty} \sum_{\substack{\rho \\ |\gamma| < T_k}} \frac{x^{\rho}}{\rho \zeta'(\rho)} - 2 + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n)! \, n \zeta(2n+1)} (\frac{2\pi}{x})^{2n}$$

Where  $\zeta(z)$  is the Riemann zeta function, and  $\rho$  are all the all nontrivial zeros of the Riemann zeta function and  $M_0(x)$  is defined as  $M_0(x) = M(x) - \frac{1}{2}\mu(x)$  if  $x \in Z^+$ , M(x) Otherwise (Odlyzko and te Riele<sup>1</sup>)

### 2. NEW PROPERTIES OF MERTENS FUNCTION

Lehman<sup>5</sup> proved that  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor) = 1$ . In general,  $\sum_{i=1}^{x} M(\lfloor x/(in) \rfloor) = 1$ , n = 1, 2, 3, ..., x(since  $\lfloor \lfloor x/n \rfloor/i \rfloor = (\lfloor x/(in) \rfloor)$ ). Let R' denote a square matrix where element (i, j) equals 1 if j divides i or 0 otherwise. (In a Redheffer matrix, element (i, j) equals 1 if i divides j or if j = 1. Redheffer proved that the determinant of the matrix equals the Mertens Function M(x).) Let T denote the matrix obtained from R' by element-by-element multiplication of the columns by  $M\left(\left\lfloor \frac{x}{1} \right\rfloor\right), M\left(\left\lfloor \frac{x}{2} \right\rfloor\right), M\left(\left\lfloor \frac{x}{3} \right\rfloor\right)...M\left(\left\lfloor \frac{x}{3} \right\rfloor\right)$ . For example, the T matrix for x = 12 is

**Theorem** (1):  $\sum_{i=1}^{x} M([x/i]) i = A(x)$ 

**Proof:** Let us now take  $A(x) = \sum_{i=1}^{x} \varphi(i)$  where  $\varphi$  is Euler's totient function. Let U denote the matrix obtained from T by element-by-element multiplication of the columns by  $\varphi(j)$ . The sum of the columns of U then equals A(x). Now since  $i = \sum_{d|i} \varphi(d)$  we can write  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor) i$  (the sum of the rows of U) equals A(x).

By the Schwarz inequality,  $A(x)/\sqrt{x(x+1)(2x+1)/6}$  is a lower bound of  $\sqrt{\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2}$ .  $A(x) = \sum_{i=1}^{x} \varphi(i)$  Is further simplified by Walfisz<sup>7</sup> and Hardy and Wright<sup>8</sup> as

$$A(x) = \sum_{i=1}^{x} \varphi(i) = \frac{1}{2} \sum_{k=1}^{x} \mu(k) \left| \frac{x}{k} \right| \left( 1 + \left| \frac{x}{k} \right| \right) = \frac{3}{\pi^2} x^2 + O(n(\log n)^{2/3} (\log \log n)^{4/3})$$

$$\sqrt{\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2} > \frac{A(x)}{\sqrt{\frac{x(x+1)(2x+1)}{6}}} > \frac{\frac{3}{\pi^2} x^2}{\sqrt{\frac{x(x+1)(2x+1)}{6}}} = \frac{3\sqrt{3}}{\pi} \frac{x^2}{\sqrt{x(x+1)(x+1/2)}}$$

This can be further simplified to

$$\sqrt{\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2 / x} > \frac{3\sqrt{3}}{\pi} \frac{1}{\sqrt{(1+1/x)(1+1/2x)}}$$

Taking limit of infinity on both the sides, we get

$$\lim_{x \to \infty} \sqrt{\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2 / x} > \frac{3\sqrt{3}}{\pi}$$

This shows that  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2$  at large values of x is greater than  $\frac{27}{\pi^2} x^2$ .

Let  $\Lambda(i)$  denote the Mangoldt function  $(\Lambda(i) \text{ equals } log(p) \text{ if } i = p^m \text{ for some prime } p \text{ and some } m \geq 1 \text{ or } 0 \text{ otherwise})$ . Mertens<sup>9</sup> proved that  $\sum_{i=1}^x M(\lfloor x/i \rfloor) log i = \psi(x)$  where  $\psi(x)$  denotes the second Chebyshev function  $(\psi(x) = \sum_{i \leq x} \Lambda(i))$ .

**Theorem (2)**: 
$$\sum_{i=1}^{x} M\left(\left|\frac{x}{i}\right|\right) \log(i) \sigma_0(i)/2 = \log(x!)$$

**Proof:** Let  $\sigma_x(i)$  denote the sum of positive divisors function  $(\sigma_x(i) = \sum_{d|i} d^x)$ . Replacing  $\varphi(j)$  with  $\log(j)$  in the U matrix gives a similar result.

Let  $\lambda(n)$  denote the Liouville function  $(\lambda(1) = 1 \text{ or if } = p_1^{a_1}...p_k^{a_k}, \lambda(n) = (-1)^{a_1+...+a_k}$ ).  $\sum_{d|n} \lambda(d)$  Equals 1 if n is a perfect square or 0 otherwise. Let  $(x) = \sum_{n \leq x} \lambda(d)$ . Let us also assume  $H(x) = \sum_{n \leq x} \mu(n)\log(n)$ .  $H(x)/(x\log(x)) \to 0$  as  $x \to \infty$  and  $\lim_{x \to \infty} M(x)/(x - H(x)/(x\log(x))) = 0$ . The statement  $\lim_{x \to \infty} M(x)/(x - H(x)/(x\log(x))) = 0$  is equivalent to the prime number theorem. Also,  $\Lambda(n) = -\sum_{d|n} \mu(d)\log(d)$ . (Apostol<sup>10</sup>).

The generalization of the Euler's totient function is Jordan totient function. Let it be denoted as  $J_k(x)$  which is defined as number of set of k positive integers which are all less than or equal to n that will form a co-prime set of (k + 1) positive integers

together with n. Let us define  $(x) = \sum_{i=1}^{x} J_k(i)$ . It is known that  $\sum_{d|n} J_k(d) = n^k$ . Then we get the following theorem.

Theorem (3): 
$$\sum_{i=1}^{x} M\left(\left|\frac{x}{i}\right|\right) i^k = B(x)$$

B(x) is expanded by McCarthy<sup>11</sup> to be

$$B(x) = \sum_{i=1}^{x} J_k(i) = \frac{n^{r+1}}{(r+1)\zeta(r+1)} + O(n^r)$$

We therefore get 
$$\sum_{i=1}^{x} M\left(\left\lfloor \frac{x}{i} \right\rfloor\right) i^k = B(x) = \frac{n^{k+1}}{(k+1)\zeta(k+1)} + O(n^k)$$
.

Likewise we can derive some other similar relationships using the T matrix that are as listed below:

**Theorem (4)**: 
$$\sum_{i=1}^{x} M\left(\left|\frac{x}{i}\right|\right) \sigma_k(i) = \sum_{i=1}^{x} i^k$$
 for  $k \in \mathbb{Z}^+$ 

**Theorem (5):**  $\sum_{i=1}^{x} M\left(\left\lfloor \frac{x}{i} \right\rfloor\right)$  where the summation is over those *i* values that are perfect squares equals L(x)

Theorem (6): 
$$\sum_{i=1}^{x} M\left(\left|\frac{x}{i}\right|\right) \Lambda(i) = -H(x)$$

**Theorem** (7): 
$$\sum_{i=1}^{x} M\left(\left\lfloor \frac{x}{i} \right\rfloor\right) 2^{\omega(n)} = \sum_{i=1}^{x} |\mu(i)| \sim \frac{6}{\pi^2} x^2 + O(\sqrt{n}) = No. of Square Free Integers$$

No. of Square Free Integers

**Theorem (8):**  $\sum_{i=1}^{x} M\left(\left\lfloor \frac{x}{i} \right\rfloor\right) d(n^2) = \sum_{i=1}^{x} 2^{\omega(i)}$  where d(x) is the sum of all the divisors of x

Theorem (9): 
$$\sum_{i=1}^{x} M\left(\left|\frac{x}{i}\right|\right) d^2(n) = \sum_{i=1}^{x} d(i^2)$$

**Theorem (10):** 
$$\sum_{i=1}^{x} M\left(\left|\frac{x}{i}\right|\right) \left(\frac{i}{\varphi(i)}\right) = \sum_{i=1}^{x} \frac{\mu^{2}(i)}{\varphi(i)}$$

Similarly many other relationships can be found between various arithmetic functions and the Mertens Functions.


## 3. A LIKELY UPPER BOUND OF |M(x)|


The following conjecture is based on data collected for  $x \le 500,000$ .

**Conjecture** (1): 
$$log(x!) > \sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2 > \psi(x)$$
 when  $x > 7$ 

By Stirling's formula, log(x!) = x log(x) - x + O(log(x)), since log(x) increases more slowly than any positive power of log(x), this is a better upper bound of  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2$  than  $x^{1+\varepsilon}$  for any  $\varepsilon > 0$ . This likely bound can be used to prove the Riemann Hypothesis since  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor) > |M(x)|$  and therefore we can write  $\sqrt{\log(x!)} > |M(x)|$ . Since the growth of  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2$  is lesser than  $x^{1+\varepsilon}$  for any  $\varepsilon > 0$ . We can say


$$M(x)x^{-\frac{1}{2}-\epsilon} \to 0 \text{ as } x \to \infty.$$






**Fig 1:** Plot of  $log(x!), \sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2$ , and  $\psi(x)$  for x = 1, 2, 3, ..., 1000

Let  $j(x) = \sum_{i=1}^x M(\lfloor x/i \rfloor)^2$  where the summation is over i values where i|x. Let  $l_1, l_2, l_3$  denote the x values where j(x) is a local maximum (that is, greater than all preceding j(x) values) and let  $m_1, m_2, m_3$ .... denote the values of the local maxima. The local maxima occur at x values that equal products of powers of small primes (Lagarias<sup>12</sup> discussed colossally abundant numbers and their relationship to the Riemann hypothesis). See Figure 2 for a plot of  $l_i/(log(l_i) m_i)$ ,  $m_i/l_i$ , and  $1/log(l_i)$  for  $i=1,2,3,\ldots,772$  (corresponding to the local maxima for  $x \le 15,000,000,000$ ). (M(x) Values for large x were computed using Del'eglise and Rivat's<sup>13</sup> algorithm.) The first two curves cross frequently, so there are i values where mi is approximately equal to  $l_i/\sqrt{log(l_i)}$ .




**Fig 2:** Plot of  $l_i/(log(l_i) m_i)$ ,  $m_i/l_i$ , and  $1/log(l_i)$  for i = 1, 2, 3, ..., 772


See Figure 3 for a plot of j(x) and  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2$  for x=1, 2, 3, ..., 10,000. See Figure 4 for a plot of  $log(l_i), log(m_i), log(M(l_i)^2)$ , and  $log(m_i/\sigma_0(l_i))$  for i=1, 2, 3, ..., 772 (when M(li)=0, log(M(li) 2) is set to -1). See Figure 5 for a plot of  $|M(l_i)|/\sqrt{l_i}$  for i=1,2,3,...,772. The largest known value of  $|M(x)|/\sqrt{x}$  (computed by Kotnik and van de Lune<sup>14</sup> for  $x \le 1014$ ) is 0.570591 (for M(7,766,842,813)=50,286). The largest  $|M(l_i)|/\sqrt{l_i}$  value for  $x \le 15,000,000,000$  is 0.568887 (for  $l_i=7,766,892,000$ ). The largest known value of  $|M(x)|/\sqrt{x}$  (computed by Kuznetsov<sup>15</sup> is 0.585767684 (for M(11,609,864,264,058,592,345)=-1,995,900,927).)



**Fig 3:** Plot of j(x) and  $\sum_{i=1}^{x} M(\lfloor x/i \rfloor)^2$  for x = 1, 2, 3, ..., 10,000




**Fig 4:** Plot of  $log(l_i)$ ,  $log(m_i)$ ,  $log(M(l_i)^2)$ , and  $log(m_i/\sigma_0(l_i))$  for i = 1, 2, 3, ..., 772



**Fig 5:** Plot of  $|M(l_i)|/\sqrt{l_i}$  for i = 1, 2, 3, ..., 772

Let  $l_i$  and  $m_i$  be similarly defined for the function  $\sigma_0(x)$ . ( $l_i$ , i=1,2,3,... are known as "highly composite" numbers. Ramanujan<sup>16</sup> initiated the study of such numbers. Robin<sup>17</sup> computed the first 5000 highly composite numbers.) Let  $m_i'$  denote  $j(l_i)$ . See Figure 6 for a plot of  $l_i/(log(l_i)m_i')$ ,  $m_i'/l_i$ , and  $1/log(l_i)$  for i=2,3,4,...,160 (corresponding to the local maxima for  $x \le 2,244,031,211,966,544,000$ ). (M(x) values for large x were computed using an algorithm similar to that used by Kuznetsov. The computations were done on an Intel i7-6700K CPU with 64 GB of RAM.) Although the first two curves cross frequently,  $m_i'$  does not appear to converge to  $l_i/\sqrt{log(l_i)}$ .



**Fig 6:** Plot of  $l_i/(log(l_i)m_i')$ ,  $m_i'/l_i$ , and  $1/log(l_i)$  for i = 2, 3, 4, ..., 160






Fig 7: Plot of  $log(l_i) + log(log(l_i)), log(l_i), log(m'_i)$ , and  $log(M(l_i)^2)$ , for i = 2, 3, 4, ..., 160


See Figure 7 for a plot of  $log(l_i) + log(log(l_i)), log(l_i), log(m'_i),$  and  $log(M(l_i)^2)$ , for i=2,3,4,...,160 (when  $M(l_i)=0, log(M(l_i)^2)$ , is set to -1). The vertical distance between the first and third curves appears to become roughly constant. See Figure 8 for a plot of  $(log(l_i) + log(log(l_i)) - log(m'_i),$  for i=2,3,4... 160. See Figure 9 for a plot of  $log(l_i) + \frac{1}{2}log(log(l_i)), log(\sum_{i=1}^{l_i} M([l_i/i])^2)$ , and  $log(l_i)$  for i=2,3,4,..., 160.  $log(l_i) + \frac{1}{2}log(log(l_i))$  Is greater than  $log(\sum_{n=1}^{l_i} M([l_i/n])^2)$  and  $log(\sum_{n=1}^{l_i} M([l_i/n])^2)$  is greater than  $log(l_i)$  for i>4. This is evidence in support of Conjecture 1. See Figure 10 for a plot of  $log(l_i) + \frac{1}{2}log(log(l_i)) - log(\sum_{n=1}^{l_i} M([l_i/n])^2)$  for i=2,3,4,..., 160.



**Fig: 8** for a plot of  $(log(l_i) + log(log(l_i)) - log(m'_i)$ , for i = 2, 3, 4... 160.



**Fig 9:** Plot of  $log(l_i) + \frac{1}{2}log(log(l_i))$ ,  $log(\sum_{i=1}^{l_i} M([l_i/i])^2)$ , and  $log(l_i)$  for i = 2, 3, 4, ..., 160



**Fig 10:** Plot of  $log(l_i) + \frac{1}{2}log(log(l_i)) - log(\sum_{n=1}^{l_i} M(\lfloor l_i/n \rfloor)^2)$  for i = 2, 3, 4, ..., 160.

#### 4. CONCLUSION

In this paper we derived new relations between Mertens function with a different arithmetic functions and also discussed about a likely upper bound of the absolute value of the Mertens function  $\sqrt{\log(x!)} > |M(x)|$  when x > 1 with sufficient numerical evidence. More experimental evidence can be found in the work of Cox and Ghosh<sup>18-19</sup>. Using this likely upper bound we showed that we have a sufficient condition to prove the Riemann Hypothesis using the Littlewood condition.

#### REFERENCES

- [1] Odlyzko, A. M. and te Riele, H. J. J. "Disproof of the Mertens Conjecture." J. reine angew. Math. 357, 138-160, 1985.
- [2] Stieltjes, T. C. R. A. S. 1885.
- [3] J. E. Littlewood, Quelques cons\_equences de l'hypoth\_ese que la fonction\_(s) n'a pas de z\_eros dans le demi-plan Re(s) > 1=2. C. R. Acad. Sci. Paris 154, 263-266 (1912)
- [4] Titchmarsh, E. C. The Theory of Functions, 2nd ed. Oxford, England: Oxford University Press, 1960.
- [5] R. S. Lehman, On Liouville's Function, Math. Comput. 14:311-320 (1960)
- [6] R. M. Redheffer, Eine explizit losbare Optimierungsaufgabe, Internat. Schriftenreine Numer. Math., 36 (1977)
- [7] Walfisz, A. Ch. 5 in Weyl's exponential sums in modern number theory. Berlin: Deutscher Verlag der Wissenschaften, 1963.
- [8] Hardy, G.H. and Wright, E.M. "The Average Order of phi (n)." §18.5 in An Introduction to the Theory of Numbers, 5th ed. Oxford, England: Clarendon Press, pp. 268-269, 1979
- [9] F. Mertens, Uber eine zahlentheoretische Funktion, Akademie Wissenschaftlicher Wien Mathematik-Naturlich Kleine Sitzungsber, 106 (1897) 761-830
- [10] T. M. Apostol, Introduction to Analytic Number Theory, Springer, 1976
- [11] McCarthy, P.J.: Introduction to Arithmetical Functions. Springer, New York (1986)
- [12] J. C. Lagarias, An elementary problem equivalent to the Riemann hypothesis, American Mathematical Monthly 109, 534-543 (2002)
- [13] M. Del\_eglise and J. Rivat, Computing the Summation of the Mobius Function, Experimental Mathematics 5, 291-295 (1996)
- [14] T. Kotnik and J. van de Lune, On the order of the Mertens function, Experimental Mathematics 13, pp. 473-481 (2004)
- [15] E. Kuznetsov, Computing the Mertens function on a GPU, arXiv:1108.0135v1 [math.NT], (2011)

- [16] S. Ramanujan, Highly composite numbers, Proc. London Math. Soc. 14 (1915), 347-407
- [17] G. Robin, M\_ethodes d'optimalisation pour un probl\_eme de th\_eories des nombres., RAIRO Inform. Th\_eor. 17, 239-247, 1983
- [18] Cox, Darrell & Ghosh, Sourangshu. (2021). Signs of the Mertens function. 10.13140/RG.2.2.26074.47042.
- [19] Cox, Darrell & Sultanow, Eldar & Ghosh, Sourangshu. (2021). The Energy Spectral Density of the Mertens Function. 10.13140/RG.2.2.33037.67043.